

New species of *Leptothorax* (Hymenoptera: Formicidae) from Cuba

Nuevas especies de *Leptothorax* (Hymenoptera: Formicidae) de Cuba

Jorge Luis Fontenla Rizo

Museo Nacional de Historia Natural de Cuba, Obispo 61, 10100, La Habana, Cuba

Abstract

Five new species of *Leptothorax* from Cuba are described: *L. abeli*, from Sierra Maestra, Santiago de Cuba Province, arboreal habits and related to *L. darlingtoni* y *L. splendens*; *L. banoao*, from Sancti Spiritus Heights, Sancti Spiritus Province; limestone dweller species and related to *L. myersi* and *L. villarensis*; *L. cuyaguatete*, from Northern Serpentine Heights, Pinar del Río Province, terricolous habits with no defined relationships; *L. imias*, from Imías, Santiago de Cuba Province, terricolous habits with no defined relationships, and *L. nipensis*, from Nipe-Sagua-Baracoa Massif, limestone dweller species, related to *L. mortoni* and *L. porphyriti*.

Resumen

Se describen cinco especies nuevas de *Leptothorax* de Cuba, *L. abeli*, de la Sierra Maestra, Provincia de Santiago de Cuba, de hábitos arborícolas y relacionada con *L. darlingtoni* y *L. splendens*; *L. banoao*, de las alturas de Sancti Spiritus, de formaciones cársicas, relacionada con *L. myersi* y *L. villarensis*; *L. cuyaguatete*, de las alturas pizarrosas del Norte, Provincia de Pinar del Río, de hábitos terrícolas y sin afinidades precisas; *L. imias*, de Imías, Provincia de Santiago de Cuba, de hábitos terrícolas y sin afinidades precisas, y *L. nipensis*, de las alturas de Nipe-Sagua-Baracoa, Provincia de Holguín, de formaciones cársicas relacionada con *L. mortoni* y *L. porphyriti*.

Key words: *Leptothorax*, new species, Formicidae, Cuba.

Palabras clave: *Leptothorax*, especies nuevas, Formicidae, Cuba.

INTRODUCTION

Leptotracine are the most remarkable group of Cuban ants, because its diversity (more than 30 described species), endemism (only three species are not exclusive of Cuba), and the heterogeneous arrangement of morphological and ecological features, as a consequence of the evolutionary radiation that the group has developed mostly in Cuba territory. BARONI-URBANI (1978) revised the Caribbean segment of the genus *Leptothorax*, formerly considered as *Macromischa*, and described several new species, including Cuban. He also defined 12 morphological species groups. Later, SNELLING (1986) and FONTENLA (1997) synonymized some of the species previously recognized in Baroni Urbani's revision. I am currently undertaking a phylogenetic analysis of the Antillean *Leptothorax*. As a result, some of the Baroni-Urbani's morphological species groups will change in definition and composition. Because of this reason, the species described in this paper won't be formally included in none of those groups.

All of the specimens are deposited at the Museo Nacional de Historia Natural de Cuba.

Measurements and indices.

Morphological terminology in the text follows BOLTON (1994).

Measurements (mm). Scape length (SL), head width (HW), mesosoma length (ML), petiole length (PL), petiole height (PH), postpetiole width (PPW), postpetiole length (PPL). Both of the postpetiole measures in dorsal view. Posterior femur length (FL). Indices. Scape index (SI). SL/HL X100; cephalic index (CI): HW/HL X100; petiole index (PI): PL/MLX100; petiole height index (HI): PH/PLX100. Postpetiole index (PPI): PPW/PPLX100; femur index (FI): FL/MLX100.

SYSTEMATIC

Leptothorax abeli new species (Fig. 1)

Measurements. Workers (paratypes) SL: 1,12 (1,12-1,2), HW: 1,2 (1,1-1,2), HL: 1,4 (1,4-1,42), ML: 2,1 (1,9-2,1), PL: 0,78 (0,70-0,78), PH: 0,55 (0,52-0,55), PPW: 0,6 (0,6-0,53), PPL: 0,37 (0,37-0,40), FL: 1,55 (1,150-1,55). Indices. SI: 80,3 (80,3-85,7), CI: 82,1 (73,8-82,1), PI: 36,9 (33,3-36,9), HI: 70,9 (70,9-75), PPI: 160 (131-160), FI: 73,8 (73,8-79,4).

Material examined. Holotype: worker, collected by Abel Pérez on May 20, 1995, in Subida a la Nigua, Sierra Maestra, Santiago de Cuba Province. Paratypes: 8 workers, same data as holotype; 4 workers, collected by A. G. Debras, on July 25, in Rio Peladeros, Sierra Maestra, Santiago de Cuba Province.

Diagnosis and comparison. Arboreal species of Cuban *Leptothorax* with a very distinctive pattern of polichrous coloration and heavy mesosoma rugae, except in the anterior dorsal part of pronotum. Short propodeal spines present, tibiae and femora swollen with very small tubercles. Petiolar node scale-shaped and postpetiole broader than long. This species is related to *L. darlingtoni* (Wheeler) and *L. splendens* (Mann). It differs from both of them by lacking the heavy rugae in the anterior part of pronotum in dorsal view. It is more similar to *L. splendens* in general size and body proportions, but differs in the more vivid color pattern, broader petiolar node and smoother head. Both species have not overlapping distribution (*L. abeli* in Western Sierra Maestra, *L. splendens* in the Nipe-Sagua-Baracoa Masiff).

Description. Mandibles five toothed, slightly striated. Antennae with 12 segments, club well differentiated and three-segmented; funiculus segments rounded and stout; scapes short and stout, no surpassing the cephalic length. Head subquadrated. Mesosoma massive, curved in profile, almost evenly broad in dorsal view, throughout. Humeral angles rounded. Promesonotal depression not very conspicuous. Propodeal spines short (16,6% mesosoma length) and stout but sharp at the tips, slightly curved outward in the middle and divergent at the tips; propodeum with a gently declive. Petiolar peduncle short, with 36,9% of mesosoma length, petiolar node scale-shaped and high (54,8% of petiole length). Postpetiole broader than longer. Gaster with a well developed sting. Femora and

tibiae swollen and with small scattered tubercles; femora do not surpass mesosoma length. Tegument shining throughout. Head surface polished, with only feeble striae on the fron and surrounded the eyes. Mesosoma with thick sulcae, transverse at the dorsum and diagonal on the pleurae. Dorsal anterior part of pronotum lacks the sulcae, surface smooth, only with very faint longitudinal striae. Body pilosity composed by long and white acuminate hairs, mostly erected, abundant, but not densely packed. Colour of head, antennae, legs, petiole, postpetiole and gaster shining brown. Smooth part of pronotum with violaceous reflections. Rest of mesosoma darker brown with conspicuous greenish reflections. Coxae yellowish brown.

Distribution. Apparently distributed by the western part of the Sierra Maestra.

Etimology. Named after Abel Pérez, a Cuban arachnologist and first collector of the species.

Comments. The specimens were observed in relatively high numbers, running on tree trunks and branches. Caught specimens reacted in aggressive way, stinging painfully.

Leptothorax banao new species

(Fig. 2)

Measurements Workers (paratypes). SL: 1.62 (1.56-1.62), HW: 0.99 (0.93-0.96), HL: 1.23 (1.20-1.23), ML: 1.83 (1.80-1.83), PL: 1.2 (1.1-1.2), PH: 0.33 (0.30-0.33), PPW: 0.39 (0.36-0.39), PPL: 0.42 (0.39-0.42), FL: 1.86 (1.83-1.86). Indices. SI: 131.7 (130-131.7), CI: 80.5 (78.0-80.5), PI: 65.6 (60.1-65.6), HI: 27.5 (27.2-27.5), PPI: 92.8 (92.3-92.8), FI: 101.8 (101.6-101.8).

Material examined. Holotype: worker, collected by Abel Pérez on May, 1994, in Banao, Sancti Spiritus Heights, Sancti Spiritus Province. Paratypes: 4 workers, same data as holotype.

Diagnosis and comparison. Limestone-dweller species of Cuban *Leptothorax*, with a long and slender body and elongated scapes, legs, petiole and propodeal spines, petiolar node low and rounded, postpetiole longer than broad. Color reddish brown, gaster brown, small and shining. This species is related to *L. villarensis* and *L. myersi*. It is more similar to *L. villarensis*, with similar low petiolar nodes, but it differs in minor size, remarkable more slender constitution and lower petiolar node. Besides, coloration is deeper, pilosity more scarce, and longitudinal striation pattern of dorsum of mesosoma is more homogeneous. In addition, it differs from *L. myersi* in the low petiolar node, which is very high and almost scale shape in this last species. *L. banao*, so far it is known, lives in geographic isolation in the Sancti Spiritus Heights, separated by the *Agabama* basin from the distributional range of the other species, which are sympatric in the Trinidad Heights and some surrounded lowlands.

Description. Mandibles five toothed. Funiculum with 11 segments. Antennal club four-segmented, but little differentiated; scapes very long and slender, surpassing cephalic length. Eyes relatively small. General constitution of the body slender, mesosoma with longitudinal profile; promesotoracic depression pronounced; propodeal spines very




Fig. 1. A, *Leptothorax abeli* new sp. 625X (tegumentary sculpture and most of pilosity no depicted). B, *Leptothorax banao* new sp. 625X (tegumentary sculpture and most of pilosity no depicted). C, *Leptothorax cuyaguateje* new sp. 625X (tegumentary sculpture and most of pilosity no depicted). D, *Leptothorax imias* new sp. 625X (tegumentary sculpture no depicted). E, *Leptothorax imias* female. 312.5X (tegumentary sculpture no depicted). F, *Leptothorax nipensis* new sp. 625X (tegumentary sculpture and most of pilosity no depicted).

Fig. 1. A, *Leptothorax abeli* esp. nov. 625X (esculturación y mayor parte de la pilosidad no reflejada). B, *Leptothorax banao* esp. nov. 625X (esculturación y mayor parte de la pilosidad no reflejada). C, *Leptothorax cuyaguateje*, esp. nov. 625X (esculturación y mayor parte de la pilosidad no reflejada). D, *Leptothorax imias* esp. nov. 625X (esculturación no reflejada). E, *Leptothorax imias* hembra. 312.5X (esculturación no reflejada). F, *Leptothorax nipensis* esp. nov. 625X (esculturación y mayor parte de la pilosidad no reflejada).

long (almost 40% of mesosoma length) and sharp, divergent basally, curved outward in the middle and convergent at the tips. Petiole very long and slender (more than 60% of mesosoma length), node very low and rounded, with no defined faces; postpetiole bell-shaped, longer than broad. Gaster proportionally small, sting well developed. Femora long, surpassing mesosoma length; medially portion not abruptly swollen; tibiae normal. Tegument shining in the gaster; head surface with longitudinal narrow striae; mesosoma sides and dorsum of pronotum with longitudinal rugae; rest of mesosoma dorsum with transverse rugae. Petiole striated; postpetiole slightly striated. Pilosity abundant, except in petiole and postpetiole, composed by long white acuminate hairs. Colour reddish; antennae and legs brown; gaster black.

Distribution. Sancti Spiritus Heights, Sancti Spiritus Province.

Etimology. Named after the type locality.

Comments. The specimens were collected on limestone rocks in a forest habitat. It is assumed this species is a limestone-nesting, not only because the habitat, but also because the high resemblance to *L. myersi* and *L. villarensis*, which built the typical tubular cartoon-like entrance to the nest, located in limestone rocks crevices. Also the specimens displayed the characteristic slow movements of this kind of species.

Leptothorax cuyaguateje new species
(Fig. 3)

Measurements. Workers (paratypes) SL: 0.66 (0.64-0.66), HW: 0.54 (0.52-0.54), HL: 0.72 (0.68-0.72), ML: 0.81 (0.79-0.81), PL: 0.33 (0.30-0.33), PH: 0.27 (0.25-0.27), PPW: 0.39 (0.37-0.39), PPL: 0.30 (0.25-0.30), FL: 0.57 (0.55-0.57). Indices. SI: 91.7 (91.7-94.1), CI: 75.0 (75.0-76.5), PI: 40.7 (37.9-40.7), HI: 81.8 (81.8-83.3), PPI: 130.0 (130.0-148.0), FI: 70.4 (69.6-70.4).

Material examined. Holotype, worker, from Ceja de Francisco, Northern Serpentine Heights, Pinar del Rio Province, collected by J. L. Fontenla in March, 1993. Paratypes; three workers, same data as holotype.

Diagnosis and comparison. Terricolous species of Cuban *Leptothorax* with small size, massive constitution, convex profile, long propodeal spines, petiolar node scale-shaped, very broad postpetiole, femora swollen, pilosity scarce with short and blunt hairs, and polymorphic coloration. This species shows a general resemblance with the species of the heterogeneous "*pulchellus*" Baroni Urbani's group. All of these species share some traits like terricolous habits, scarce pilosity composed by short and blunt hairs and scapes and posterior femora not surpassing cephalic and mesosoma length respectively. *L. cuyaguateje* does not seem obviously close to any of them in particular, according to the combination of traits listed in the diagnosis.

Description. Mandibles five toothed. Funiculum with 11 segments, antennal club well differentiated and three segmented, scapes slender, not surpassing cephalic length. Me-

mesosoma robust, with convex profile, and promesotoracic suture no deep. Propodeal spines long, almost 30% of mesosoma length, straight and divergent from base to tips. Petiole long, about 40% of mesosoma length. Petiolar node scale-shaped and high. Postpetiole bell-shaped, broader than long. Gaster relatively big. Femora short, not surpassing mesosoma length and swollen. Tegument slightly shining. Head, mesosoma, petiole and postpetiole punctuated; petiolar node and gaster smooth. Pilosity scarce, composed by short and blunt white erected hairs. Antennae and legs with more abundant hairs, very short and appressed. Coloration variable, dark brown throughout, or opaque yellowish in head and mesosoma, with the remaining body areas dark brown.

Distribution. Only known from the type locality.

Etimology. Aborigine voice. It is the name of the largest river in the Pinar del Rio Province.

Comments. The specimens were collected in pine tree forest litter, growing in serpentine soil, very close to the typical “mogotes” hills of the Sierra de los Organos. It is the first known species of Antillean *Leptothorax* associated at this particular habitat.

Leptothorax imias new species
(Fig. 4)

Measurements. Workers (paratypes). SL: 0.53 (0.50-0.53); HW: 0.57 (0.55-0.57). HL: 0.66 (0.62-0.65); ML: 0.90 (0.85-0.90); PL: 0.42 (0.40-0.42); PPW: 0.18 (0.15-0.18); PPL: 0.27 (0.22-0.27); FL: 0.60 (0.55-0.60). Indices. SI: 80.3 (80.3-80.6); CI: 86.4 (86.4-88.7); PI: 46.7 (44.4-46.7); HI: 42.8 (37.5-42.8); PPI: 133.3 (133.3-136.3); FI: 66.7 (64.7-66.7).

Material examined. Holotype; worker, collected by L. F. de Armas in August, 1975, Imias, Santiago de Cuba Province. Paratypes: four workers, one female, same data as holotype.

Diagnosis and comparison. Terricolous species of Cuban *Leptothorax* with small size, yellow color, longitudinal profile, propodeal spines short, femora swollen, no hairs on the dorsum of mesosoma and very broad and flattened postpetiole. This species shares the common traits already mentioned above with terricolous Antillean *Leptothorax*. Its affinities are not obvious, and the combination of traits like lacking hairs in the dorsum of mesosoma and a very broad and flattened postpetiole make this species very distinctive among the terricolous group.

Description. Mandibles five toothed. Antennae with 12 segments; club conspicuous and three segmented; scapes slender, not surpassing cephalic length. Head and eyes proportionally big. Mesosoma robust with longitudinal profile. In dorsal view, pronotum with humeral angles rounded; mesosoma width decreasing from pronotum to propodeum. Promesotoracic suture superficial. Propodeal spines short, about 17% of mesosoma length, straight and divergent at the tips. Petiolar peduncle short, node high, almost square in profile and rounded in dorsal view. Postpetiole very broad and flattened dorso-

ventrally. Gaster proportionally big, with a well developed sting. Femora swollen and short, posterior femora less than 70% of mesosoma length. Tegument shining; body punctuated through and very feeble striated. Gaster smooth. Pilosity extremely scarce, with only a few short and blunt white hairs restricted mostly to gaster and clypeus; dorsum of mesosoma with no hairs. Color yellow brown. Mandibles with dentarium border red.

Female. (Fig. 5). Measurements. SL: 0.60; HW: 0.69; HL: 0.87; ML: 1.29; PL: 0.51; PH: 0.27; PPW: 0.57; PPL: 0.36; FL: 0.72. Indices. SI: 68.9; CI: 79.3; PI: 39.5; HI: 52.9; PPI: 52.6; FI: 55.8.

Description. In comparison with workers, size much bigger, antennal club less conspicuous, eyes very large. Head with three ocelli, mesosoma very robust; petiolar node rounded and massive, postpetiole proportionally small. Tegument, pilosity and coloration like the workers.

Distribution. Only known from the type locality.

Etimology. Named after the type locality.

Comments. The specimens were collected from a very small nest under a stone. Imias is a locality with a very dry climate and xerophitic vegetation.

Leptothorax nipensis new species
(Fig. 6)

Measurements. Workers (paratypes). SL: 1.35 (1.30-1.35); HW: 0.87 (0.84-0.90); HL: 1.26 (1.23-1.29); ML: 1.83 (1.75-1.83); PL: 0.93 (0.90-0.93); PH: 0.30 (0.27-0.30); PPW: 0.36 (0.33-0.36); PPL: 0.45 (0.42-0.45); FL: 1.83 (1.77-1.83). Indices. SI: 107.1 (104.6-107.7); CI: 69.1 (68.3-69.8); PI: 52.6 (51.4-52.6); HI: 32.3 (30.0-32.4); PPI: 80.0 (78.6-80.0); FI: 102.0 (100.0-102.0).

Material examined. Five workers, collected by A. R. Estrada on July 1994, at the limestone cutter Julio Antonio Mella, South of Sierra de Nipe, Santiago de Cuba Province.

Diagnosis and comparison. Limestone-dweller species of Cuban *Leptothorax*, which builds a cartoon tubular entrance to the nest. Body slender with very elongated scapes and legs, femora slightly swollen, tibiae normal, propodeal spines very long, petiole long with a low and rounded node. Postpetiole longer than broad. Color brown reddish. This species is related to *L. mortoni* from the South of Santiago de Cuba and Guantánamo Provinces and to *L. porphyritis*, from the Habana-Matanzas Heights. It is easily distinguished from *L. mortoni*, because lacks the so distinctive color pattern of the gaster, has more shining tegument, more slender body and appendages and the propodeal spines are more curved in the middle. Both species are alopatric; *L. mortoni* is associated with coastal habitat. It differs from *L. porphyritis* in the more slender body and appendages, better defined longitudinal striated pattern over a densely micropunctuated surface, denser pilosity and tegument less shining.

Description. Mandibles five toothed. Antennae with 12 segments; club little differentiated; scapes slender, surpassing cephalic length; eyes relatively small. Mesosoma with longitudinal profile and slender; pronotum and metanotum broad in comparison with mesonotum, which is relatively constricted. Humeral angles rounded and promesotoraxical suture conspicuous. Propodeal declive very low; propodeal spines long and sharp (36% of mesosoma length); divergent at the base, curved in the middle and straight at the tips. Petiole very long, more than half the mesosoma length, petiolar node rounded and very low, with no defined faces. Postpetiole longer than broad, slender. Gaster proportionally small, with a well developed sting. Posterior femora length similar to mesosoma length, not very swollen; tibiae normal and long. Tegument slightly shining throughout, densely punctuated and feeble but well defined striae. Gaster only with very small punctures. Pilosity dense, white hairs long, and acuminate. Body color dark reddish brown; legs, spines, petiole and postpetiole with lighter tone.

Etimology. Named derived from Sierra de Nipe.

Distribution. Only known from the type locality.

Comments. *L. nipensis* was observed walking slowly in sunny hours on the surface of limestone walls, quite naked of vegetation and exposed to presumed predators. This behaviour is very similar to the other limestone-dweller species of Cuban *Leptothorax*.

ACKNOWLEDGMENTS

To A. Pérez and A. R. Estrada for the collect of specimens, and to N. García for help in the sketches.

BIBLIOGRAPHY

- BARONI-URBANI, C. 1978. Materiale per una revisione dei *Leptothorax* Neotropicali appartenente al sottogenere *Macromischa* Roger, n. comb. (Hymenoptera: Formicidae). *Entomol. Basil.*, 3: 395-618.
- BOLTON, B. 1994. *Identification guide to the ant genera of the world*. Harvard University Press, Cambridge, Massachusetts.
- FONTENLA, J. L. 1997. Notas y sinónimos nuevos de *Leptothorax* (Hymenoptera: Formicidae) de Cuba. *Avicennia*, 6/7: 47-53.
- SNELLING, R. 1986. New synonymy in Caribbean ants of the genus *Leptothorax* (Hymenoptera: Formicidae). *Proc. Ent. Soc. Washington*, 88: 154-156.